

**CONFIGURATIONS OF RANK- $40r$ EXTREMAL EVEN
UNIMODULAR LATTICES ($r = 1, 2, 3$)**

SCOTT DUKE KOMINERS AND ZACHARY ABEL

ABSTRACT. We show that if L is an extremal even unimodular lattice of rank $40r$ with $r = 1, 2, 3$ then L is generated by its vectors of norms $4r$ and $4r+2$. Our result is an extension of Ozeki's analogous result for the case $r = 1$.

1. INTRODUCTION

A *lattice of rank n* is a free \mathbb{Z} -module of rank n equipped with a positive-definite inner product $(\cdot, \cdot) : L \times L \rightarrow \mathbb{R}$. The *dual* of L , denoted L^* , is the set

$$L^* = \{y \in L \otimes \mathbb{R} : \forall x \in L, (x, y) \in \mathbb{Z}\},$$

which itself forms a lattice of the same rank as L . For a lattice vector $x \in L$, we call (x, x) the *norm* of x . A lattice L is *integral* if $(x, x') \in \mathbb{Z}$ for all $x, x' \in L$, i.e. if and only if $L \subseteq L^*$. An integral lattice is said to be *unimodular* if it is self-dual ($L = L^*$).

A lattice L is called *even* if and only if every lattice vector has an even integer norm, i.e. $(x, x) \in 2\mathbb{Z}$ for $x \in L$. An even lattice is automatically integral by the familiar parallelogram identity, $2(x, x') = (x + x', x + x') - (x, x) - (x', x')$.

Lattices that are simultaneously even and unimodular are especially rare. Indeed, such a lattice's rank must be divisible by 8. Sloane proved that if L is an even unimodular lattice of rank n then the minimal (nonzero) norm in L is bounded by

$$(1) \quad \min_{\substack{x \in L \\ x \neq 0}} (x, x) \leq 2\lfloor n/24 \rfloor + 2$$

(see [2, p. 194, Cor. 21]). An even unimodular lattice of rank n is called *extremal* if it attains the bound (1).

Ozeki [6, 8] showed that if L is an extremal even unimodular lattice of rank 32 or 48 then L is generated by its vectors of minimal norm. The first author [5] showed analogous results for extremal even unimodular lattices of ranks 56, 72, and 96. In a similar vein, Ozeki [7] showed that if L is extremal even unimodular of rank 40, then L is generated by its vectors of norms 4 and 6. Here, we extend and slightly simplify Ozeki's methods, recovering Ozeki's rank-40 result and obtaining analogous results for extremal even unimodular lattices of ranks 80 and 120.

Date: December 5, 2007.

2000 Mathematics Subject Classification. 11H06 (52C07, 05B30).

Key words and phrases. Even unimodular lattices, extremal lattices, weighted theta series.

2. MODULAR FORMS AND THETA SERIES

We will use the notation $\mathcal{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ for the *upper half plane* of complex numbers. A *modular form of weight k for the group $PSL_2(\mathbb{Z})$* is a holomorphic function $f : \mathcal{H} \rightarrow \mathbb{C}$ which is holomorphic at $i\infty$ and satisfies

$$f\left(\frac{az+b}{cz+d}\right) = (cz+d)^k f(z)$$

for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PSL_2(\mathbb{Z})$. If a modular form f vanishes at $z = i\infty$, it is called a *cusp form*.

Let M_k and M_k^0 be the \mathbb{C} -vector spaces of modular forms and cusp forms of weight k respectively. It is known that the *Eisenstein series*

$$\begin{aligned} E_4(z) &= 1 + 240e^{2\pi iz} + 2160e^{4\pi iz} + 6720e^{6\pi iz} + \dots \text{ and} \\ E_6(z) &= 1 - 504e^{2\pi iz} - 16632e^{4\pi iz} - 122976e^{6\pi iz} - \dots, \end{aligned}$$

which are modular forms of weights 4 and 6 respectively, freely generate the spaces M_k in the sense that any nonzero modular form can be written uniquely as a weighted homogeneous polynomial in E_4 and E_6 . This implies that $\dim(M_k) = 0$ for k odd, negative, or $k = 2$; that $\dim(M_{2k}) = 1$ and $\dim(M_{2k}^0) = 0$ for $k = 0$, $2 \leq k \leq 5$ and $k = 7$; and that multiplication by the weight-12 modular form $\Delta = 12^{-3}(E_4^3 - E_6^2)$ defines an isomorphism $M_{k-12} \xrightarrow{\sim} M_k^0$. More information on the theory of modular forms for $PSL_2(\mathbb{Z})$ can be found in [9].

The *theta function* $\Theta_L : \mathcal{H} \rightarrow \mathbb{C}$ associated to a lattice L is defined by

$$\Theta_L(z) = \sum_{x \in L} e^{\pi i(x,x)z};$$

it is a generating function encoding the norms of L 's vectors. For a homogeneous *harmonic polynomial* $P \in \mathbb{C}[x_1, \dots, x_n]$, i.e. a homogeneous polynomial for which $\sum_{j=1}^n \frac{\partial^2 P}{\partial x_j^2} = 0$, we define the *weighted theta series* $\Theta_{L,P}$ by

$$\Theta_{L,P}(z) = \sum_{x \in L} P(x) e^{\pi i(x,x)z}.$$

As shown in [9, 3], if L is an even unimodular lattice of rank n then Θ_L is a modular form of weight $\frac{n}{2}$, and if in addition P is a homogeneous harmonic polynomial of degree d , then $\Theta_{L,P}$ is a modular form of weight $\frac{n}{2} + d$.

3. MAIN RESULT

We denote by $P_{d,x_0}(x)$ the “zonal spherical harmonic polynomial” of degree d , related to the *Gegenbauer polynomial* by

$$(2) \quad P_{d,x_0}(x) = G_d((x, x_0), ((x, x)(x_0, x_0))^{1/2}),$$

where $G_d(\cdot, \cdot)$ is the homogeneous polynomial of degree d such that $G_d(t, 1)$ is the Gegenbauer polynomial of degree d evaluated at t [1].

We let L be an extremal even unimodular lattice of rank $40r$, and adopt the notation used by Ozeki in [7]: For an even unimodular lattice L , we denote by $\Lambda_{2m}(L)$ the set of vectors in L having norm $2m$. We denote by $\mathcal{L}_{2m}(L)$ the sublattice of L generated by $\Lambda_{2m}(L)$, and similarly denote by $\mathcal{L}_{2m_1+2m_2}(L)$ the sublattice of L generated by $\Lambda_{2m_1}(L) \cup \Lambda_{2m_2}(L)$.

We define $a(2k, L) := |\Lambda_{2k}(L)|$. It is clear that the theta series Θ_L is given by $\Theta_L(z) = \sum_{k=0}^{\infty} a(2k, L)e^{2k\pi iz}$. We note that

$$4r = 2\lfloor 5r/3 \rfloor + 2 = \min\{2k > 0 : a(2k, L) \neq 0\}$$

is the minimal norm of vectors in L and use the notation

$$\begin{aligned} N_j(x) &= |\{y \in \Lambda_{4r}(L) : (x, y) = j\}|, \\ M_j(x) &= |\{y \in \Lambda_{4r+2}(L) : (x, y) = j\}|. \end{aligned}$$

Using the involution $y \longleftrightarrow -y$ of $\Lambda_m(L)$, we see that we have $N_j(x) = N_{-j}(x)$ and $M_j(x) = M_{-j}(x)$ for any $j \in \mathbb{R}$ and $x \in L \otimes \mathbb{R}$.

We will show the following configuration result, which directly extends Ozeki's [7] result for extremal even unimodular lattices of rank 40:

Theorem 3.1. *For $r = 1, 2, 3$ and L extremal even unimodular of rank $40r$, we have $L = \mathcal{L}_{4r+(4r+2)}(L)$.*

Proof. We partition L into its equivalence classes modulo $\mathcal{L}_{4r+(4r+2)}(L)$. We need only show that any class $[x] \in L/\mathcal{L}_{4r+(4r+2)}(L)$ is represented by a vector $x_0 \in [x]$ with $(x_0, x_0) \leq 4r + 2$.

Now, we suppose there exists some equivalence class $[x_0] \in L/\mathcal{L}_{4r+(4r+2)}(L)$ where $x_0 \neq 0$ is a representative of minimal norm with $(x_0, x_0) = 2t$ for some $t \geq 2r + 2$. We have the inequality

$$(3) \quad |(x_0, x)| \leq 2r \text{ for all } x \in \Lambda_{4r}(L),$$

as x_0 is not minimal in L whenever $(x_0, \pm x) > 2r$ since the vector $x \mp x_0$ has norm

$$(x \mp x_0, x \mp x_0) = (x, x) \mp 2(x, x_0) + (x_0, x_0) < (x_0, x_0).$$

Similarly, we have

$$(4) \quad |(x_0, x)| \leq 2r + 1 \text{ for all } x \in \Lambda_{4r+2}(L).$$

From (3) and (4), we have the equations

$$(5) \quad \sum_{x \in \Lambda_{4r}(L)} (x, x_0)^{2k} = \sum_{j=1}^{2r} 2 \cdot j^{2k} \cdot N_j(x_0),$$

$$(6) \quad \sum_{x \in \Lambda_{4r+2}(L)} (x, x_0)^{2k} = \sum_{j=1}^{2r+1} 2 \cdot j^{2k} \cdot M_j(x_0),$$

for all $k > 0$.

We extract from the theta series Θ_L of L the coefficients $a(4r, L)$ and $a(4r+2, L)$. We observe immediately from (5) and (6) that

$$(7) \quad \sum_{x \in \Lambda_{4r}(L)} (x, x_0)^0 = a(4r, L),$$

$$(8) \quad \sum_{x \in \Lambda_{4r+2}(L)} (x, x_0)^0 = a(4r+2, L).$$

Since L is even unimodular of rank $40r$, we have $\Theta_{L, P_d, x_0} \in M_{20r+d}^0$ for any $d > 0$. By comparing power-series coefficients, we then observe

$$(9) \quad \Theta_{L, P_d, x_0} \equiv 0 \text{ for } d \in \{2, \dots, 4r-2, 4r+2\},$$

$$(10) \quad \Theta_{L, P_{4r}, x_0} \equiv c_1 \Delta^{2r} \text{ for a constant } c_1,$$

$$(11) \quad \Theta_{L, P_{4r+4}, x_0} \equiv c_2 E_4 \Delta^{2r} \text{ for a constant } c_2.$$

From (9), we obtain the equations

$$(12) \quad \sum_{x \in \Lambda_{4r}(L)} (x, x_0)^{2d} = a(4r, L) \frac{1 \cdot 3 \cdots (2d-1)}{40r \cdot (40r+2) \cdots (40r+2d-2)} (8r)^d t^d \quad \text{and}$$

$$(13) \quad \sum_{x \in \Lambda_{4r+2}(L)} (x, x_0)^{2d} = a(4r+2, L) \frac{1 \cdot 3 \cdots (2d-1)}{40r \cdot (40r+2) \cdots (40r+2d-2)} (8r+4)^d t^d,$$

for $d \in \{2, \dots, 4r-2, 4r+2\}$. We obtain from (10)

$$(14) \quad \sum_{x \in \Lambda_{4r+2}(L)} P_{4r, x_0}(x) = c_{4r} \sum_{x \in \Lambda_{4r}(L)} P_{4r, x_0}(x),$$

where $\Delta^{4r} = e^{(4r)\pi iz} + c_{4r} e^{(4r+1)\pi iz} + O(e^{(4r+2)\pi iz})$. Similarly, (11) gives

$$(15) \quad \sum_{x \in \Lambda_{4r+2}(L)} P_{4r+4, x_0}(x) = c_{4r+4} \sum_{x \in \Lambda_{4r}(L)} P_{4r+4, x_0}(x),$$

where $E_4 \Delta^{4r} = e^{(4r)\pi iz} + c_{4r+4} e^{(4r+1)\pi iz} + O(e^{(4r+2)\pi iz})$.

Combining the equations (7), (8), (12), (13), (14), and (15) with (5) and (6), we obtain a system of $4r+4$ homogeneous linear equations in the $4r+3$ unknowns

$$N_0(x_0), \dots, N_{2r}(x_0), M_0(x_0), \dots, M_{2r+1}(x_0).$$

At this stage, we diverge from our natural generalization of Ozeki's original methods and obtain the (extended) determinants of these inhomogeneous linear systems; these determinants must vanish because the system is overdetermined.

For $r = 1, 2, 3$, these determinants are respectively

$$(16) \quad 2^{55} 3^7 5^8 7^4 11^4 13^1 19^6 23^3 \cdot (t-2) \cdot t \cdot (6t-13) \cdot (10t^2 - 55t + 77),$$

$$(17) \quad 2^{132} 3^{27} 5^{16} 7^{10} 11^6 13^{10} 23^4 41^8 43^6 47^3 \cdot (t-4) \cdot t \cdot Q_2(t),$$

$$(18) \quad 2^{244} 3^{48} 5^{26} 7^{13} 11^7 13^7 17^6 23^4 31^{11} 37^1 59^{14} 61^{11} 67^5 71^3 73^1 \cdot (t-6) \cdot t \cdot Q_3(t),$$

where $Q_2(t)$ is the irreducible quintic

$$10768t^5 - 242280t^4 + 2202310t^3 - 10101795t^2 + 23361877t - 21771246$$

and $Q_3(t)$ is the irreducible septic

$$\begin{aligned} & 19989882674056909935t^7 - 892881426107875310430t^6 \\ & + 17258039601222654151533t^5 - 187053310321121904306075t^4 \\ & + 1227398249908229181423784t^3 - 4874010945909263810320032t^2 \\ & + 10840974078436271024624064t - 10414527769923133690990080. \end{aligned}$$

In each case, there are no integer solutions $t \geq 2r + 2$. However, we had assumed the existence of an equivalence class

$$[x_0] \in L/\mathcal{L}_{4r+(4r+2)}(L)$$

with minimal-norm representative $x_0 \neq 0$ having $(x_0, x_0) = 2t$ for integral $t \geq 2r+2$; since no such t exists, all equivalence classes must be generated by vectors having norms $4r$ and $4r + 2$. \square

4. CONCLUDING REMARKS

A quick inspection will show that our results are the only possible immediate extensions of Ozeki's methods. In the cases $r \geq 4$, it is not possible to extract sufficiently many linear conditions by these exact techniques, as the dimensions of the relevant spaces of cusp forms grows too large.

However, using different analysis, Elkies [4] has shown a stronger result than our Theorem 3.1 in the $r = 3$ case: If L is an extremal unimodular lattice of rank 120 then $L = \mathcal{L}_{12}(L)$. This result for rank-120 lattices is analogous to Ozeki's [6, 8] results in dimensions 32 and 48, and to the first author's [5] results in dimensions 56, 72, and 96.

ACKNOWLEDGEMENTS

The authors are especially grateful to Professor Noam D. Elkies, under whose direction this research was conducted. They appreciate his instruction, support, and commentary on drafts of this article. The authors also thank Denis Auroux, Katherine Paur, and Erin Schlumpf for their translation assistance and an anonymous referee for especially helpful comments and advice.

REFERENCES

- [1] C. BACHOC, G. NEBE, B. VENKOV, *Odd unimodular lattices of minimum 4*. Acta Arithmetica **101** (2002), 151–158.
- [2] J. H. CONWAY, N. J. A. SLOANE, *Sphere Packing, Lattices and Groups* (3rd edition). Springer-Verlag, New York, 1999.
- [3] W. EBELING, *Lattices and Codes* (2nd edition). Vieweg, Germany, 2002.
- [4] N. D. ELKIES, *On the quotient of an extremal Type II lattice of rank 40, 80, or 120 by the span of its minimal vectors*. Preprint.
- [5] S. D. KOMINERS, *Configurations of extremal even unimodular lattices*. To appear, Int. J. Num. Thy. (Preprint arXiv:0706.3082, 21 Jun 2007.)
- [6] M. OZEKI, *On even unimodular positive definite quadratic lattices of rank 32*. Math. Z. **191** (1986), 283–291.
- [7] M. OZEKI, *On the structure of even unimodular extremal lattices of rank 40*. Rocky Mtn. J. Math. **19** (1989), 847–862.
- [8] M. OZEKI, *On the configurations of even unimodular lattices of rank 48*. Arch. Math. **46** (1986), 247–287.
- [9] J.-P. SERRE, *A Course in Arithmetic*. Springer-Verlag, New York, 1973.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY

E-mail address: SDK: kominers@fas.harvard.edu ZA: zabel@fas.harvard.edu